

Measuring the halo mass of Mg II absorbers from their cross-correlation with Luminous Red Galaxies

Nicolas Bouché¹, M. T. Murphy², C. Péroux³ and I. Csabai⁴

¹Max Plank Institut für extraterrestrische Physik, Giessenbachstrasse, Garching D-85748, Germany; nbouche@mpe.mpg.de

²Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK;

³European Southern Observatory, Karl-Schwarzschild-Str 2, D-85748 Garching, Germany;

⁴Dept. of Physics, Eötvös Loránd University, Budapest, Pf. 32, H-1518 Budapest, Hungary.

Abstract. We study the cross-correlation between 716 Mg II quasar absorption systems and $\sim 100,000$ Luminous Red Galaxies (LRGs) selected from the Sloan Digital Sky Survey Data Release 3 in the redshift range $0.4 \leq z \leq 0.8$. The Mg II systems were selected to have $\lambda\lambda 2796$ & 2803 rest-frame equivalent widths $\geq 1.0 \text{ \AA}$ and identifications confirmed by the Fe II $\lambda 2600$ or Mg I $\lambda 2852$ lines. Over co-moving scales $0.2\text{--}13h^{-1} \text{ Mpc}$, the Mg II–LRG cross-correlation has an amplitude 0.69 ± 0.09 times that of the LRG–LRG auto-correlation. Since LRGs have halo-masses of $10^{13} M_{\odot}$, this strong cross-correlation signal implies that the absorber host-galaxies have halo-masses $1\text{--}2 \times 10^{12} M_{\odot}$.

Keywords. cosmology: observations — galaxies: evolution, halos — quasars: absorption lines

1. Introduction

The connection between quasar (QSO) absorption line (QAL) systems and galaxies (first established by Bergeron & Boissé 1991) is important to our understanding of galaxy evolution. QALs provide detailed information about the physical conditions and kinematics of galaxies out to large impact parameters ($R > 100 \text{ kpc}$), regardless of the absorber’s intrinsic luminosity (e.g. Steidel et al. 2002, and Churchill et al. 2005, Kacprzak et al. 2005, these proceedings). Past results show that Mg II absorbers are biased towards late-type galaxies which do not evolve strongly from $z \simeq 1$ (Steidel & Sargent 1992; Steidel et al. 1994). These results also show that Mg II absorber host-galaxies have K -band luminosities consistent with normal $0.7L_B^* \text{ Sb}$ galaxies. The cross-section of Mg II absorbers with $W_r^{\text{MgII}} \geq 0.30 \text{ \AA}$ appears to be $R_{\times} \sim 70h^{-1} \text{ kpc}$ (co-moving) (e.g. Steidel 1995). These systems are associated with H I absorbers in the Lyman limit regime up to the damped Ly-alpha absorber (DLA) regime (see also Rao et al. 2005, these proceedings).

In Bouché et al. (2004), we used the Sloan Digital Sky Survey (SDSS) data release 1 (DR1; Abazajian et al. 2003) to constrain the mass of the halos associated with the Mg II absorbers. Specifically, we used the absorber-galaxy cross-correlation to measure of the mass ratio of the halos associated with Mg II since, in a hierarchical galaxy formation scenario, the amplitude ratio of the Mg II–LRG cross-correlation to the LRG–LRG auto-correlation is also their bias ratio. The reader is referred to Bouché et al. (2004) and Bouché et al. (2005a) for the details. Fig. 1 (left) illustrates the methodology. Using 212 Mg II absorbers and $\sim 20,000$ Luminous Red Galaxies (LRGs), Bouché et al. (2004)




Figure 1. *Left*: The bias $b(M)$ as a function of the halo mass M Mo & White (2002) (solid line). The right y -axis shows the auto-correlation length r_0 . LRGs have $r_{0,gg} \simeq 6 h^{-1}$ Mpc, and thus masses of $\simeq 10^{13} M_{\odot}$. *Right*: Filled circles show the Mg II–LRG cross-correlation $w_{ag}(r_{\theta})$ between 716 Mg II absorbers and 94,649 LRGs. Filled triangles show the LRG–LRG auto-correlation, w_{gg} . The dashed line shows a power-law fit to w_{gg} . The solid line shows the fit $\hat{w}_{ag} = a \times \hat{w}_{gg}$ for $r_{\theta} > 200 h^{-1}$ kpc since the smallest scales will be affected by the finite cross-section of the absorbers. The raw relative amplitude is $a = 0.87 \pm 0.08$. The left panel therefore implies that our Mg II absorbers have halos 7–10 times less massive than LRG halos, i.e. our Mg II absorbers have halos with mass $1\text{--}2 \times 10^{12} M_{\odot}$.

found that the bias ratio $b_{\text{Mg II}}/b_{\text{LRG}}$ is 0.67 ± 0.09 on scale $r_{\theta} > 200 h^{-1}$ kpc, implying a halo mass for the Mg II host galaxies of $0.5\text{--}2.5 \times 10^{12} M_{\odot}$ (for $10^{13} M_{\odot}$ LRG halos).

2. Results

Here, we extend our DR1 results using SDSS Data Release 3 (DR3; Abazajian et al. 2005). We selected 716 Mg II absorbers from SDSS/DR3 with $z_{\text{abs}} \leq 0.8$ using automated technique that included the following criteria: (i) $W_r^{\text{Mg II}} \geq 1.0 \text{ \AA}$; (ii) we require that $W_r^{\text{Mg I}} \geq 0.2 \text{ \AA}$, and that $W_r^{\text{Fe II}} \geq 0.5$ following the DLA criteria of Nestor et al. (2003) and Rao & Turnshek (2000) (see Rao et al. 2005, these proceedings, for an updated discussion). We remove spurious candidates by visually inspecting each Mg II spectrum.

For each absorber, we selected ~ 1300 Luminous Red Galaxies (LRGs) from SDSS/DR3 using color criteria following Scranton et al. (2003), and in a slice of width $W_z = 0.1$ using the photometric redshift calculated with the code of Csabai et al. (2003). There are a total of 94,649 LRGs meeting these criteria, within $12.8 h^{-1}$ Mpc, our largest bin.

For the cross-correlation, w_{ag} , we use the estimator $1 + w_{ag}(r_{\theta}) = \text{AG}/\text{AR}$, where AG is the total observed number of absorber–galaxy pairs between $r_{\theta} - dr/2$ and $r_{\theta} + dr/2$ and AR is the total absorber–random galaxy pairs. This estimator is necessary to account for the non-symmetric situation: Mg II absorbers have precise redshifts, while the LRGs have photometric redshifts with an accuracy of $\sigma_z \simeq 0.1$ (see Bouché et al. 2005a, for a discussion). Fig. 1(right) shows our results (see caption). The errors to w_{ag} and w_{gg} were computed using $N_{\text{jack}} = 10$ jackknife realizations.

The amplitude of the Mg II–LRG cross-correlation relative to that of the LRG–LRG auto-correlation is $0.69 \pm 0.07 \pm 0.06$, after applying a correction of 25 ± 10 per cent discussed in Bouché et al. (2004). The two error terms reflect the statistical and systematic uncertainty, respectively. By adding the errors in quadrature, the bias ratio is

$$a = 0.69 \pm 0.09. \quad (2.1)$$

Within the context of hierarchical galaxy formation, Eq. 2.1 implies that our Mg II absorbers have halo masses 7–10 times smaller than the LRGs. For $10^{13} M_{\odot}$ LRG halos, the Mg II absorbers have halos of $1\text{--}2 \times 10^{12} M_{\odot}$.

It is important to realize that this method (i.e. measuring the halo mass from the ratio of projected correlation functions) has the following advantages (see also Bouché et al. 2004): (i) it constrains the mass of the Mg II/DLA host-galaxies in a statistical manner without directly identifying them. (ii) it is free of systematics from contaminants (e.g. stars), and (iii) it does not require knowledge of the true width of the redshift distribution of the galaxy used. The last two points are a consequence of the fact that we use the same galaxies for $w_{\text{gg}}(r_{\theta})$ and for $w_{\text{ag}}(r_{\theta})$.

3. Discussion

Our results are consistent with those of Bergeron & Boissé (1991), Mo & Miralda-Escudé (1996): For instance, Mo & Miralda-Escudé (1996) indicate that the majority of Mg II absorbers reside in systems with $V_{\text{circ}} = 150\text{--}300 \text{ km s}^{-1}$ with a median at $\sim 200 \text{ km s}^{-1}$. Our mass measurement appears to corroborate that of Steidel et al. (1994) who found that Mg II absorbers with $W_{\text{r}}^{\text{MgII}} \geq 0.3 \text{ \AA}$ are associated with late-type $\sim 0.7 L_B^*$ galaxies, since the expected amplitude ratio between early and late type galaxies is ~ 0.70 (see Bouché et al. 2004).

Are our results consistent with Λ CDM? That is, is there enough massive halos to account for dN/dz ? From $dN/dz = n(M) \sigma(M) dr/dz$, $R_X \simeq 70 \text{ kpc}$ (co-moving) (Steidel 1995), $n(M) = 10^{-2} h^{-3} \text{ Mpc}^{-3}$, $dN/dz = 0.3 (n/10^{-2}) (R_X/70 \text{ kpc})^2 \simeq dN/dz(\text{obs})$, and we can conclude that there are enough massive $10^{12} M_{\odot}$ halos. While we defer a more detailed analysis of these results to Bouché et al. (2005b), preliminary results also indicate little dependence of the halo mass on the equivalent width.

Acknowledgements

N.B. thanks the IAU and the organizers of this conference for the travel grant. M.T.M. thanks PPARC and I.C. thanks OTKA, T047244. Funding for the Sloan Digital Sky Survey (SDSS) has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Aeronautics and Space Administration, the National Science Foundation, the U.S. Department of Energy, the Japanese Monbukagakusho, and the Max Planck Society.

References

- Abazajian K. et al., 2003, AJ, 126, 2081
- Abazajian K. et al., 2005, AJ, 129, 1755
- Bergeron J., Boissé P., 1991, A&A, 243, 344
- Bouché N., Murphy M. T., Péroux C., 2004, MNRAS, 354, 25L
- Bouché N. et al., 2005a, ApJ, in press, astro-ph/0504172
- Bouché N., Murphy M. T., Péroux C., Csabai I., 2005b, MNRAS, in prep.
- Csabai I. et al., 2003, AJ, 125, 580
- Mo H. J., Miralda-Escudé J., 1996, ApJ, 469, 589
- Mo H. J., White S. D. M., 2002, MNRAS, 336, 112
- Nestor D. B., Rao S. M., Turnshek D. A., Vanden Berk D., 2003, ApJ, 595, L5
- Rao S. M., Turnshek D. A., 2000, ApJS, 130, 1
- Scranton R. et al., 2003, Phys. Rev. Lett., submitted, preprint (astro-ph/0307335)
- Steidel C. C., Sargent W. L. W., 1992, ApJS, 80, 1
- Steidel C. C., Dickinson M., Persson S. E., 1994, ApJ, 437, L75

- Steidel C. C., 1995, in Meylan G., ed., QSO Absorption Lines. Springer-Verlag, Berlin, Germany, p. 139
- Steidel C. C., Kollmeier J. A. et al., 2002, ApJ, 570, 526